

ANTIBIOGRAM

2010

NUMBERS ARE PERCENT SUSCEPTIBLE

GRAM POSITIVE ORGANISMS	No. tested	Ampicillin	Augmentin	Cefazolin	Cefepime	Ceftriaxone	Cefuroxime	Ciprofloxacin	Clindamycin-NU	Doxycycline	Erythromycin-NU	Gentamicin	Gent-AMP Synergy	Linezolid	Nitrofurantoin-U	Oxacillin	Penicillin	Rifampin	Synercid	Trim-Sulfa	Vancomycin	
<i>Staphylococcus aureus</i> (MSSA)	874		99	99		97		89	82		68	98		100		100 ^α	20	100.0		99	100	
<i>Staphylococcus aureus</i> (MRSA)	353		0	0		0		29	74		16	95		100		0 ^α	0	99.5		95	100	
<i>Staphylococcus aureus</i> -CF	202		86	86		84		78	87		49	96		99.5		86	17	94.0		99.5	100	
<i>Staphylococcus</i> spp. (coagulase negative)	400		31	31		30		57	27		27	60		100	99	32	7	96		44	100	
<i>Streptococcus pyogenes</i>	102					100			95		95						100				100	
<i>Streptococcus pneumoniae</i> ^β	69					86 ^β	78 ^β		79		75						84 ^β				72	100
Viridans streptococci ^γ	44				68	70			77		27						32					100
<i>Enterococcus faecium</i> (VRE only)	33	0								38				92				25	100			0
<i>Enterococcus</i> spp. (un-specified)	133	89						26					91	81	81							100

MRSA: methicillin resistant *Staphylococcus aureus*. MRSA rate for 2010: 29%.

MSSA: methicillin susceptible *Staphylococcus aureus*.

Percent NONSUSCEPTIBLE includes:

α: Methicillin resistance was tested also by latex agglutination of the PBP and/or molecular assay for *mecA* gene.

β: 12% ceftriaxone MIC are in the 'INTERMEDIATE' range

β: 3% cefuroxime MIC are in the 'INTERMEDIATE' range

β: 13% penicillin MIC are in the 'INTERMEDIATE' range

γ: 9% cefepime MIC are in the 'INTERMEDIATE' range

γ: 9% ceftriaxone MIC are in the 'INTERMEDIATE' range

γ: 57% penicillin MIC are in the 'INTERMEDIATE' range

U: tested on urinary isolates

NU: tested on non-urinary isolates

Seattle Children's Hospital

4800 Sand Point Way N.E. PO Box 5371 MS: A6901

Seattle, Washington 98105

Xuan Qin, Ph.D. Director (206) 987-2586

Microbiology Laboratory: (206) 987-2585

© 2011 Seattle Children's Hospital

All Rights Reserved