

Wāween Lale iaļ in Kōto ilo lien Idiņ (Juon Ajri Eruj, im Ewōr M̄wijn̄wijn ilo Jikin bōk Kōto ñan Jipaņ an Emmenono (Tracheostomy))


Ajri ro ilo m̄oko m̄weier me elōñ aer m̄wijn̄wijn ñan jipaņ aer emmenono, ekkā aer loe apaņ ilo aer emmenono ñe m̄wijn̄wijn eo epenjak kōn kaļo (mucous) ak ej wōtlōk kein jipaņ emmenono eo .

Ajri eo ej kowaļok kakōļle in apaņ ilo an emmenono

- Eļapļok jermal ilo an emmenono, joņan m̄okaj in an emmenono, im kilen an menono
- Enana kalor in kilen, edikļok joņan oxygen ko
- Enana ak ejjab m̄oto ubōn
- Ewōr ainikien wejel im ikijjelok jān m̄wijn̄wijn eo
- Eļapļok ainikien jān turin tube in m̄wijn̄wijn eo
- Kumkum/aabḡōḡ

Kein jermal eaorōk ej aikuj in pād ippān ajri eo nājūm aolep iien.

- Tube in m̄wijn̄wijn eo me joņan ej āinwōt joņan tube eo an nājūm
- Tube in m̄wijn̄wijn eo me ej juon wōt jaij dikļok jān tube eo an nājūm (trach in idiņ)
- Nitōļ in wā (ñan tube ko wōt ewōr aer kakkobaba) kein kajjir (lubricant)
- Kein jermal in kein jorom im m̄weuiuk ko
- Pāak in bar kōmour, mejen iturin māj, mejen stoma
- Oxygen



Elaññe ilo jabdewōt iien ajri eo nājūm enaaj jako an ruj, jinoe CPR

Ewōr jipaņ ko ikijjien ukok jān Seattle Children's im eļeļok wōḡnean ñan ri-nañinmej ro Ri-jaroñron, epen aer ronjake ak rejaje Kajin Palle, ro uwaan baamle, im ri-pinej-jenkwan ikijjien kien. Seattle Children's enaaj kōḡmmane bwe mejele kein ren pojak ilo wāween ko jet ne kwōj kajjitoķ. Kūri Jikin Jipaņ eo an Baamle ilo 206-987-2201.

Em̄oj an ri-jermal in Seattle Children's etale peba in leto-letak in. Bōtaab, aikuj ko an ajri eo nājūm rej an make wōt. Mokka jān am̄ kōḡmman jabdewōt ak atare mejele kein, jouj im kōnono ippān an ajri eo nājūm ri-leļok kōjparok. © 2018 Seattle Children's, Seattle, Washington. Aolep mejele ko ijin rej ñan kein jermal
© 2018 Seattle Children's, Seattle, Washington. Em̄oj dāpij aolep maroņ ko.

Emergency Airway Management (Conscious Child with Tracheostomy)


Kids at home with trachs can often have trouble breathing when the trach becomes blocked with mucus or the trach comes out.

Child is showing signs of respiratory distress

- Increased work of breathing, respiratory rate, retractions
- Poor color, drop in oxygen levels
- Poor or no chest rise
- Whistling or wheezy sounds from trach
- Increased vocalization around trach tube
- Agitated/upset

Always have essential equipment with your child at all times

- Tracheostomy tube the same size as your child's
- Trach tube one size smaller (emergency trach)
- Syringe (for cuffed tubes only) and lubricant
- Suction equipment and supplies
- Resuscitation bag, face mask, stoma mask
- Oxygen



If at any time your child loses consciousness, start CPR.

Seattle Children's offers interpreter services for Deaf, hard of hearing or non-English speaking patients, family members and legal representatives free of charge. Seattle Children's will make this information available in alternate formats upon request. Call the Family Resource Center at 206-987-2201.

This handbook has been reviewed by clinical staff at Seattle Children's. However, your child's needs are unique. Before you act or rely upon this information, please talk with your child's healthcare provider.

© 2018 Seattle Children's, Seattle, Washington. All rights reserved.