

A Guide to Feminizing Hormones

Gender Affirming Care

Hormone therapy is an option that can help transgender people feel more comfortable in their bodies. Like other medical treatments, there are benefits and risks. Knowing what to expect will help us partner to maximize the benefits and minimize the risks.

The binary term “male,” “female,” “masculine,” “feminine,” “masculinizing” and “feminizing” do not accurately reflect the diversity of people’s bodies or identities. To describe how hormones work, it is helpful to know how testosterone works in non-intersex, non-trans men’s bodies, and how estrogen and progesterone works in non-intersex, non-trans women’s bodies. We keep these binary terms in quotes to emphasize that they are artificial and imperfect concepts.

What are hormones?

Hormones are chemical messengers that tell tissues of the body how to function, when to grow, when to divide and when to die. They regulate many functions, including growth, sex drive, hunger, thirst, digestion, metabolism, fat burning and storage, blood sugar, cholesterol levels and reproduction.

What are sex hormones?

Sex hormones are involved in the development of the penis and testicles, or the vulva and clitoris (external genitals). Sex hormones also affect the secondary sex characteristics that typically develop at puberty (facial and body hair, bone growth, breast growth, voice changes, etc.). There are 3 categories of sex hormones in the body:

- **Androgens:** testosterone, dehydroepiandrosterone (DHEA), dihydrotestosterone (DHT)
- **Estrogens:** estradiol, estriol, estrone
- **Progestin:** progesterone

Generally, people with testicles tend to have higher androgen levels, and people with ovaries tend to have higher levels of estrogens and progestogens.

What is hormone therapy?

Hormone therapy is taking medicine to change the levels of sex hormones in your body. Changing these levels will affect your hair growth, voice pitch, fat distribution, muscle mass and other features that are associated with sex and gender. Feminizing hormone therapy can help make the body look and feel less “masculine” and more “feminine” — making your body more closely match your identity.

To Learn More

- Adolescent Medicine - 206-987-2028
- Gender Clinic Care Navigator
206-987-8319
- Ask your child’s healthcare provider
- seattlechildrens.org

Free Interpreter Services

- In the hospital, ask your nurse.
- From outside the hospital, call the toll-free Family Interpreting Line, 1-866-583-1527. Tell the interpreter the name or extension you need.

What medicines are involved?

There are different kinds of medicines used to change the levels of sex hormones in your body. These medicines work by affecting:

- The part of your brain that stimulates sex hormone production
- Your testicles (which produce testosterone)
- The cells in your body that respond to sex hormones

Usually, feminizing hormone therapy involves:

- Estrogen
- A medicine to block testosterone
- A combination of estrogen and a medicine to block testosterone
- Sometimes a progestin is added

Estrogen

Estrogen is the main hormone responsible for promoting “feminine” physical traits. It works directly on tissues in your body (for example, makes breasts develop). Estrogen also indirectly reduces testosterone. Estrogen can be taken by:

- Pill (oral or under the tongue)
- Injection (intramuscular or subcutaneous)
- Skin patch or gel (transdermal)

There are different formulations of estrogen. Your healthcare provider will talk to you about the different kinds and what is right for you.

Androgen blockers

Androgen-blockers work by blocking testosterone. They are also known as anti-androgens or androgen antagonists. They reduce “masculine” physical traits and have a mildly “feminizing” effect. For example, they will help slow “male” pattern baldness, reduce growth of facial hair and stop spontaneous/morning erections.

There are different types of androgen blockers. The one most typically prescribed is spironolactone. Androgen blockers are often prescribed in addition to estrogen because they have effects that complement each other. Taking androgen blockers reduces the amount of estrogen you need to get the same effects, which minimizes the health risks associated with estrogen. Androgen blockers can be prescribed alone for people who want to reduce “masculine” characteristics for a more androgynous appearance because it is less “feminizing” than estrogen.

Progestins

There are mixed opinions about using progestins for feminizing hormone therapy. Some gender clinic programs choose not to use progestins due to the lack of clear evidence that they facilitate “feminization.” Progestins also have known side effects (which include depression, weight gain and changes to blood fats).

However, progestins may be used by some gender care providers in the following situations:

- If estrogen alone is not working, even at the maximum dose.
- As a replacement for estrogen if there are concerns about the side effects or risks of estrogen.
- To promote nipple and breast development (but there is not strong evidence for this yet).

As with estrogen and androgen blockers, balancing possible risks and benefits of progestins is a decision between you and your healthcare providers.

What is a typical dose?

Feminizing hormone therapy varies greatly from person to person. There is no right hormone combination, type, or dose for everyone. Deciding what to take depends on your health because each hormone therapy has different risks and side effects. What your healthcare provider prescribes depends on what is available locally and what you can afford. It may also depend on insurance coverage.

It can change based on how your body reacts when you start taking hormones — everyone's body is different and sometimes people have a negative reaction to a specific kind of medicine.

The right dose or type of medicine for you might not be the same as for someone else. It is a good idea to discuss the advantages and disadvantages of different options with us. If you have any concerns about being able to take the medicines — or about the side effects, costs or health risks — let us know. We take your needs and concerns into account when planning your hormone therapy.

We will partner with you to explore your insurance coverage and any other resources to make sure you get the care you need.

In prescribing a specific medicine and dose, we consider your overall health, including any other medicines you are taking. Every person is different — each body absorbs, processes and responds to sex hormones differently. Some people show more changes than others. Changes happen more quickly for some than others.

Taking more hormones than the dose you were prescribed will not speed up changes. Taking more than your prescribed dose greatly increases your health risks.

If you think your dose is too low, talk with us to discuss your options. It might be better to try a different type of medicine or a combination of medicines, rather than increasing the dose.

If you have your testicles removed, your body will only produce a tiny amount of testosterone, so:

- The dose of estrogen can be reduced
- Androgen blockers can be reduced or stopped

You will need to stay on estrogen or another form of medicine for the rest of your life to keep your bones strong. We may also suggest you take low-dose testosterone to help your metabolism. Your provider may also suggest that you take calcium and vitamin D supplements to protect your bones.

What changes can I expect?

Feminizing hormone therapy has important physical and psychological benefits. Bringing mind and body closer together eases gender dysphoria and can help you feel better about your body. People who have had gender dysphoria often describe being less anxious, less depressed, calmer and happier when they start taking hormones. For some people, this psychological change happens as soon as they start taking hormones. For others, it happens a bit later as the physical changes appear more.

Each person changes differently. How quickly changes appear for you depend on:

- Your age
- The number of hormones receptors in your body
- The way your body responds to the medicine

There is no way to know how your body will respond before you start hormones.

Androgen blocker (spironolactone) without estrogen

Taking spironolactone (the most common androgen blocker) without estrogen has small effects. The changes are caused by the medicine blocking the effect of testosterone in your body. Most of the changes are reversible, which means if you stop taking it, your body will go back to how it was before you started taking the medicine. Androgen blockers affect the whole body. You cannot pick the changes you want.

Average timeline	Effect
After 1 to 3 months	<ul style="list-style-type: none">• Decreased sex drive• Fewer instances of waking up with an erection or spontaneously having an erection. Some people also have difficulty getting an erection even when they are sexually aroused• Decreased ability to make sperm and ejaculatory fluid
Gradual changes (usually takes at least 2 years)	<ul style="list-style-type: none">• Slower growth of facial or body hair• Slowed or stopped “male”-pattern balding• Slight breast growth (reversible in some cases, not in others)

Estrogen

Taking estrogen has stronger physical “feminizing” effects. These changes are caused by the estrogen’s effect on cells in your body that have estrogen receptors. Taking estrogen also has an indirect effect of suppressing testosterone production. Like androgen blockers, estrogen affect the whole body. You cannot pick the changes you want.

Average timeline	Effect
After 1 to 6 months	<ul style="list-style-type: none">• Softening of skin• Less muscle mass and more body fat• Redistribution of body fat to be more on breasts and hips• Possible decrease in sex drive• Fewer instances of waking up with an erection or spontaneously having an erection. Some people also have difficulty getting an erection even when they are sexually aroused.• Decreased ability to make sperm and ejaculatory fluid
Gradual changes (maximum changes after 2 to 3 years)	<ul style="list-style-type: none">• Nipple and breast growth• Slower growth of facial and body hair• Slowed or stopped “male” pattern balding• Smaller testicles

Breast and nipple growth starts early but is usually gradual. It can take 2 years or more for breasts to reach their maximum size. As with all people, there is a range in how large breasts grow. In many cases, your breasts might not grow beyond an A or B cup size. If you are not happy with the size of your breasts after 18 to 24 months on estrogen, you can consider surgical augmentation. The implants will look most natural if you wait to get as much growth as you can from hormones.

Most of the effects of hormones happen in the first 2 years. During this time, the doctor who prescribes your hormones will usually want to see you every 3 months. This is to check if the hormones are working properly. After that, you will probably need an appointment every 6-12 months. At appointments in the first 2 years, your doctor will likely:

- Look at your facial and body hair. If you shave, the doctor will ask how quickly your hair grows back.
- Ask about changes to your sex drive, erections, or other sexual changes.
- Ask about breast growth or nipple changes
- Order blood test to see what your hormone levels are.
- Ask how you feel about the changes that have happened so far.

After 2 years, your doctor will monitor the effects by asking if you notice any more changes from the hormones.

When you are 21 years old, you will transition to a medical provider who can continue your treatments as an adult. For information about moving to an adult healthcare provider, visit seattlechildrens.org/TransitioningToAdultHealthcare.

Are the changes permanent?

Some of the changes you will notice from the feminizing hormone therapy are not permanent. If you stop taking the medicine, some of the changes will stop and your body will return to how it was before you started the hormones. There are 3 types of changes that may be permanent:

- Breast growth
- Fertility
- Fat distribution to hips

Breast growth

If you are taking the androgen blocker called spironolactone without estrogen because you do not want visible changes, you might see some breast growth. This growth happens slowly, so you can stop taking it if you do not want breast growth. Breast growth from spironolactone is usually small and reversible. But in some people, the breast tissue remains even after the spironolactone is stopped.

Estrogen causes permanent nipple development and breast growth. Even if you stop taking estrogen, the breast tissue will not go away and your nipples will not shrink.

Fertility

Both androgen blockers and estrogen affect your production of sperm, which means you may have trouble having biological children after taking them. It is also important to know that we do not yet fully understand the long-term effects feminizing medications have on fertility. If you stop taking feminizing hormones, your ability to make sperm may or may not return to what it was before you started. We strongly recommend that you talk about options for sperm banking before starting hormone therapy. If you have already started hormones, you can work with your doctor to stop the hormones, give sperm samples and store them if they are viable. Then you could go back on hormones.

Although androgen blockers and estrogen affect sperm production, there may still be a chance you could make someone pregnant after starting hormone therapy. **Depending on how you have sex, you may need to use birth control.**

Hormone therapy does not lower your risk of HIV and other sexually transmitted infections. Depending on how you have sex, you may need to use condoms, gloves or other latex barriers. Feminizing hormones can make erections less firm, increasing the risk of condom leakage. In this situation, your partner can use a special condom they put inside their anus or vagina. They are called “female condoms,” but can be used by people of any gender.

What are the risks?

The medical effects and safety of feminizing hormone therapy are not fully understood. Most of the studies on hormone therapy involve different doses than are used for gender affirming care. There may be long-term risks that are not known yet.

We can lower many of the known risks of feminizing hormone therapy by creating a hormone combination that is made just for you. There are also actions you can take to reduce the risks, including:

- Not smoking. This is the number one thing you can do to reduce your risk of blood clots and heart disease. Even the occasional smoker is at an increased risk. If you do not smoke it increases the amount of estrogen that we can prescribe safely.
- Having your blood tested as recommended by your doctor.

Social repercussions

Being a person in a transphobic society can have social risks. Some people experience violence, harassment and discrimination, while others have lost support of loved ones. If you are worried about how others might react to the changes that come with hormone therapy, counseling can be useful. If you are looking for a therapist, see “How to Find a Therapist.” seattlechildrens.org/pdf/PE2195.pdf.

Blood clots

Taking estrogen increases the risk of blood clots. Blood clots can cause death, permanent lung damage (clot in the lungs), permanent brain damage (stroke), heart attack or chronic problems with veins in your legs. The risk of blood clots is much higher for if you smoke.

The danger is so high that some doctors will not prescribe estrogen if you smoke, even occasionally. Most healthcare providers will prescribe you only a low dose of estrogen until you fully stop smoking. The risk of blood clots can be made lower by:

- Taking estrogen by skin patch or gel (transdermal)
- Taking estrogen under the tongue (sublingual)
- Taking estrogen by injection (intramuscular or subcutaneous)
- Using a lower dose of estrogen

Taking estrogen changes the way your body uses and stores fat. Taking estrogen can increase deposits of fat around your internal organs. This type of fat is associated with an increased risk for diabetes and heart disease. Estrogen also increases the risk of gallstones, which can block your gallbladder. See a medical professional right away if you have these symptoms of gallstones:

- Chest, leg or abdominal pain
- Any swelling (edema) in your legs

If you have the following symptoms for more than a couple of days, call a healthcare professional:

- Nausea and vomiting (similar to morning sickness in pregnant women)
- Frequent headaches or migraines, if the pain is unusually bad or if you are vomiting

High blood pressure

Estrogen can also cause an increase in blood pressure. This can be avoided by taking estrogen with an androgen blocker medicine (spironolactone) that lowers blood pressure. If you cannot take spironolactone, you can make other changes to reduce your risk. This includes other types of medicine, exercise, not smoking and changes to your diet.

Galactorrhea and prolactinoma

With breast growth, there is often an increase in milky discharge from the nipples. This is called galactorrhea. This is caused by the estrogen stimulating the production of the hormone prolactin, which stimulates breast ducts to make milk. We do not know if milk production increases the risk of noncancerous tumors (prolactinoma) of the pituitary gland.

Although prolactinoma is not usually life-threatening, it can damage your vision and cause headaches. For this reason, your doctor will monitor for signs of prolactinoma regularly for at least 3 years after you start taking estrogen. More tests can be ordered if your prolactin level is high or if prolactinoma is suspected.

Breast cancer

It is not known if estrogen causes an increased risk of breast cancer. There have been cases of people who have developed breast cancer after hormone therapy for gender affirming care. Talk with your healthcare provider about screening tests that can be done to catch early signs of breast cancer. Your breast cancer risk is higher if you:

- Have a family history of breast cancer
- Have been taking estrogen or progesterin for more than 5 years
- Are 50 years or older
- Are overweight

Kidney health

Spironolactone (the most common androgen blocker) affects the balance of water and salt in the kidneys. If the amount of water and salt gets out of balance, you can have problems with low blood pressure. Rarely, this imbalance can lead to high levels of potassium in your body, which can cause changes in heart rhythm that can be life-threatening. Your blood tests will check your potassium levels and kidney function on a regular basis. This is especially important if you:

- Have a history of kidney problems
- Are taking medicine that can raise blood potassium (ask your doctor or pharmacist)
- Are taking ACE-inhibitors (commonly prescribed for people with high blood pressure or heart problems).

If you receive care from another healthcare provider, tell them you are on hormone therapy, so you do not take these kinds of medicines unknowingly.

Migraine headaches

Migraine headaches may happen more often after starting estrogen. People with a history of migraines may want to begin therapy at lower estrogen doses and increase doses slowly. Please talk with your provider if you develop new or different migraines after starting estrogen.

Skin rash

The skin patch (transdermal application) of estrogen can sometimes cause a skin rash. The androgen blocker spironolactone can also cause a skin rash. If this happens, contact us.

How do I get the most benefit and minimize risks?

You can help make hormone therapy as effective and safe as possible. Here are steps you can take:

- **Be informed.** Understanding how hormones work, what to expect, and possible side effects and risks will give you the tools to be in charge of your health and make informed decisions. Do your own research and ask questions. To get started, see “Gender Clinic Booklist and Resources” seattlechildrens.org/pdf/PE2634.pdf.
- **If you smoke, stop or cut down.** Any smoking greatly increases the risks of hormone therapy. If you are a smoker, your estrogen level may be kept low. If you need help to quit smoking, we can help you develop a plan or direct you to resources. You can contact QuitNow quitnow.net/Program/ as a first step. If you are not quite ready to quit, consider cutting down. Every little bit helps.
- **Find a healthcare provider you trust and can be honest with.** To get the most from hormone therapy, you need to be able to talk openly about what you want, concerns you have, and problems you are experiencing. You should feel comfortable to talk openly with your healthcare provider about your health history, smoking, alcohol, street drugs, dietary supplements, herbs and any other medicines you are taking. Hormone therapy can be affected by all of these things. Being honest with your healthcare provider will help the provider to create a hormone plan that is right for you.
- **Deal with problems early on.** If caught early enough, most of the problems that can result from hormone therapy can be dealt with in a creative way that does not involve stopping hormone therapy. Waiting to talk with your provider can make the problem worse.
- **Do not change medicine on your own.** Check with your healthcare provider if you want to start, stop or change the dose of any of your medicines. Taking medicine more often or at a higher dose than prescribed increases health risks and can slow down the changes you want. If you want to change your medicine, talk with your provider first.
- **Take a holistic approach to your health.** Health involves more than just hormone levels, and taking hormones is only one way for you to improve your quality of life. Building a circle of care that includes health professionals, friends, partners and other people who care about you will help you to deal with problems as they come up. This support will help you to heal from societal transphobia.
- **Know where to go for help.** The Seattle Children’s Gender Clinic can help you find information on health and transition issues. We can also help you connect with support groups and community resources. We can help with referrals if you need assistance finding other medical providers, counselors or another type of health professional.

What will not change?

Body image

Many people experience an increase in self-esteem and confidence as their body changes with hormones. You might find that there are also unrealistic societal standards after hormone therapy. It can be hard to separate gender dysphoria from body image problems. Professional and peer counseling can help you sort through your expectations about your appearance and work toward self-acceptance.

Mental health

Many people experience positive emotional changes from hormone therapy, including decreased gender dysphoria. Hormone therapy might help you to become more accepting of yourself, but life can still present emotional and social challenges. Biological factors, stresses of transphobia and unresolved personal issues can also affect your mental health. It is important to continue to access counseling, medication and other supports as needed for your mental health.

Your community

Some people hope that they will find greater acceptance after they make physical changes. Seek support from people and communities who accept and respect you as your body, gender identity and expression evolve. It can be helpful to connect with other transgender people, while remembering that no one will exactly mirror your own experience, identity and beliefs. It can be common to feel lonely and alone after starting hormone therapy. Having a support network to turn to can help.

Your body

Hormone therapy does not affect some parts of the body. Some changes are very small. Parts of the body that will not change:

- Penis
- Vagina
- Sex chromosomes
- Adam's apple
- Bone structure
- Voice pitch
- Height

Hormone therapy can make facial and body hair grow more slowly and be less noticeable, but hair will not go away completely. Some people get laser treatment or electrolysis to get rid of facial hair. Laser hair removal works best if you have light skin and dark hair. Electrolysis destroys the follicle that the hair grows out of, so it is permanent hair removal. Electrolysis works for all people.

While “male” pattern baldness may slow down or stop, bald areas will not grow hair again. Some people use wigs or hairpieces, hair transplants or other medical treatments, like Minoxidil (Rogaine).

Feminizing hormone therapy does not change how high or low your voice is (pitch). Hormone therapy will not change your speech patterns. Speech

therapy can help change pitch and other aspects of speech associated with gender. Some people have surgery on their vocal cords or the surrounding cartilage to try to make their voice sound higher.

Once your bones have stopped growing after puberty, feminizing hormone therapy cannot change the size or shape of your bones. Some people use facial feminizing surgery to change the shape of the skull and facial features, and to reduce a prominent Adam's apple. After puberty, there are no treatments you can take to change your height or the size of your hands and feet.

How often do I need to come in for appointments?

You need regular physical exams and lab tests to monitor your overall health while you are on hormone therapy. The first year after starting hormones, this will be at least every 2 to 3 months.

What will happen at appointments?

At every appointment, we will:

- Ask questions about your overall health
- Check your blood pressure, check your weight and listen to your lungs
- Look at your arms, legs, hands and feet to check your overall circulation and look for any signs of swelling, fluid retention or pain
- Check for early warning signs of health problems that can be caused by hormone therapy (blood clots, heart disease, diabetes)
- Recommend blood tests
- Recommend other tests (such as bone scans, heart stress function tests) as needed, depending on your health history, age and any signs of possible health problems
- Starting at age 40, but also depending on your age, family history and other risks for breast cancer, you may need an examination of your breast tissue (mammogram). When you are over 50, your healthcare provider should discuss checking for prostate cancer.

While gender healthcare training for providers emphasizes the need to be creative and stopping hormones only as a last resort, there are some health problems that make it dangerous to take hormones, such as uncontrolled heart disease. If your healthcare provider suspects you have one of these health problems, we will try to control it through medical treatment and changes to your diet or exercise routine. If the condition cannot be controlled, your provider may switch you to another type of hormone or reduce or stop your dose until your other health problems can be controlled.

Resources

Feminizing Hormone Therapy at Seattle Children's (video, 3:12)
youtu.be/8_gdLCXKI5Y

Excellence for Transgender Health - transhealth.ucsf.edu