

# Bladder Exstrophy v2.0: Prenatal

[Approval & Citation](#)

[Summary of Version Changes](#)

[Explanation of Evidence Ratings](#)

## PHASE I

### Inclusion Criteria

- Suspected diagnosis of bladder exstrophy

### Exclusion Criteria

- Suspected diagnosis of cloacal exstrophy

### Prenatal Consultations

- Urology
- Social Work

### Delivery Recommendations

- Delivery at any location that is comfortable with bladder exstrophy care
- Provider to provider consultation available

### Family Education

- Bladder Exstrophy: How to Care for your Child's Bladder Before Surgery [PE3009](#)
- Bladder Exstrophy [PE174](#)

### Provider Education

- [Outside Provider Goals of Care](#)

!

If child is stable,  
no need to transfer until  
mom can also be  
discharged

Go to Neonatal  
Phase

# Bladder Exstrophy v2.0: Neonatal

[Approval & Citation](#)

[Summary of Version Changes](#)

[Explanation of Evidence Ratings](#)

## PHASE II

!

Child can be cared for on the floor unless they meet standard NICU criteria

### Inclusion Criteria

- Diagnosis of bladder exstrophy

### Exclusion Criteria

- Age > 12 months
- Cloacal exstrophy

### Cloacal Exstrophy

- Will need general surgery involvement for ostomy + bladder plate closure

### Initial Management

#### Imaging

- Renal US
- Pelvic xray

#### Diet

- Regular

#### Antibiotics

- None

#### Bladder Exstrophy Care

- Tegaderm over bladder plate in hospital; Tegaderm or Press and Seal at home
- Silk tie on umbilical cord

#### Activity Restrictions

- None

#### Consultations

- Urology
- Orthopedics (including Spica tiger brought to family)
- Social work
- Lactation

#### Standard Newborn Care

!

Do not give prophylactic antibiotics prior to bladder closure

### Family Education

- Bladder Exstrophy: How to Care for your Child's Bladder Before Surgery [PE3009](#)
- Bladder Exstrophy [PE174](#)
- Latex Allergy [PE001](#)

#### Urology Folder: (Urology NP will give to family)

- Information about the Association for Bladder Exstrophy Community
- Medical Alert ID [PE2417](#)
- Urology Rounds [PE3005](#)
- Bladder Exstrophy: Care After Surgery [PE2910](#)
- Urinary Reflux [PE181](#)

### Discharge Criteria

- Teaching complete
- Consults seen
- Follow-up scheduled
- Cleared by Pediatrics

### Follow-Up Coordination

- PCP identified and appointment scheduled within 3-7 days of discharge
- Urology clinic visit in 2 months

# Bladder Exstrophy v2.0: Operative

[Approval & Citation](#)

[Summary of Version Changes](#)

[Explanation of Evidence Ratings](#)

## PHASE III

### Timing of Surgery

Elective closure at 4-6 months of age with osteotomy and spica cast

### Inclusion Criteria

- Diagnosis of bladder exstrophy

### Exclusion Criteria

- Age > 12 months
- Cloacal exstrophy

### Pre-operative Visits

2 months prior to surgery

- Urology clinic
  - Request iron deficiency anemia screen from pediatrician

2 weeks prior to surgery

- Urology clinic
- Orthopedics clinic
  - Order car seat
- PASS clinic
  - Type and Screen

### Outpatient Pre-Operative Management

- Anemia screen, if positive treat and discuss delaying surgery
- Car seat ordered at orthopedic RN visit
- OR team e-mail sent – **Urology fellow after Monday AM conference**
  - Pre-surgical Overview PowerPoint
  - Full Body Prep instructions
  - Pick lists
  - Huddle time

### Infection Control

2 weeks prior to surgery

- Bladder plate culture
  - **If positive**, treat with 7 days antibiotics to end on the day of surgery, then culture directive perioperative antibiotics
  - **If negative**, cefazolin only for perioperative antibiotics

## Intra Operative Care

### Anesthesiologist

#### Pain control:

- Epidural, TAP or iliac fascial blocks when anatomy permits
- Standing Tylenol post-operatively
- Standing Toradol post-operatively if >6 months and no renal anomalies

#### Fluids/Lines/Tubes:

- A-line and 2 large lvs
- Type and cross
- Orogastric tube. Remove at the end of case.
- mIVF on a pump with separate replacement of deficit and blood loss
- Albumin 5% as needed
- [Transfusion guide](#)

#### Infection Control:

- Antibiotics completed 30 minutes prior to incision

#### Extubation:

- Goal to extubate at end of case

### Surgical tech

- Place urinary stents in a container to collect urine
- Monitor and call out urine output for anesthesia
- Open bladder exstrophy specific pelvic osteotomy kit

### Nursing

- Bladder Exstrophy: Full body prep
- Radiopaque table
- Positioning at end of table
- **Fluoroscopy available for osteotomy**
- Page Orthopedic surgeon at end of case to place spica
- PACU nurse to petal spica

!

Fluid goal is euvolemia

Go to Post-Operative Phase

# Bladder Exstrophy v2.0: Post-Operative

[Approval & Citation](#)

[Summary of Version Changes](#)

[Explanation of Evidence Ratings](#)

## PHASE IV

### NICU Admission

- At the discretion of the surgical and anesthesia teams
- Standard criteria for NICU discharge

### Inclusion Criteria

- Diagnosis of bladder exstrophy

### Exclusion Criteria

- Age > 12 months
- Cloacal exstrophy

## Initial Management

### Family Education

- Urology Rounds [PE3005](#)

### Vital signs

- Standard vital signs per acute care guidelines
- Strict I/O

### Activity

- Out of bed ad lib
- OK to hold baby

### Fluids/Electrolytes/Nutrition

- D5 NS@maintenace IV+PO
- CBC, Chem7, Cr/BUN POD1
- Regular diet immediately after surgery

### Medications

- Pain service consult in PACU
- Oxybutynin TID standing for bladder spasms

### Wound Care

- Monitor incisions for early signs of infection

### Consults

- Lactation consultation if nursing or pumping
- Social work

**!**  
**STOP and RESOLVE if incisions are not clean, dry and intact**

## Infection Control

- Perioperative antibiotics for 24 hours after surgery
- No prophylaxis while bladder being continuously drained
- Urine culture 2 weeks prior to outpatient planned cast and tube removal
  - If positive, give culture driven antibiotics for 7 days (5 days prior to cystogram / VCUG, 2 days after)
- Antibiotic prophylaxis for vesicoureteral reflux if seen on outpatient VCUG at time of cast removal

## Family Education (once stable after surgery)

- Bladder Exstrophy [PE174](#)
- Bladder Exstrophy: Care After Surgery [PE2910](#)

## Follow-up Coordination

- Urology weekly visits x 6 weeks
- Spica cast removal at 4 weeks
  1. Orthopedics clinic visit for spica removal
    - Pelvic X-Ray to evaluate osteotomies
- 6 week visit for catheter removal
  1. Urology clinic visit for catheter removal
  2. Radiology:
    - Cystogram for bladder integrity; if (-), VCUG for VUR
    - Renal Ultrasound

## Supplies and teaching

### (care coordination to initiate)

- Appropriate car seat
- Cather flush
- Petals / Mole skin
- Diapers

## Discharge Criteria

- Teaching complete
- Supplies ordered
- Car seat approved
- Follow-up appointments scheduled
- Medically cleared by Urology and Orthopedic surgery

# Bladder Exstrophy Transfusion Guideline

## Laboratory Monitoring

- Check blood gas every hour
- Check CBC and coags after osteotomies are complete or if concern for hemodynamic instability
- Consider TXA for high risk patients (ie high blood loss expected, malnourished, history of prematurity)
- **50 mg/kg** (maximum 2000 mg) intravenous over **15** min followed by 5 mg/kg/hour continuous I.V. infusion until skin closure

### Lab Results

Hct < 21 or hemodynamic instability

Platelets < 100k

Fibrinogen < 150

*TEG > 10 or INR > 1.5*

### Intervention

PRBC 10-15 mL/kg

Platelets 5 – 10 mL/kg

Cryoprecipitate 2 - 5 mL/kg

*FFP 10-15 mL/kg*

[Return to Operative](#)

# Bladder Exstrophy Outside Provider

## Management of the bladder plate

- The goal is to keep the bladder plate clean and moist
- Tegaderm should be placed over the bladder plate overlapping with the surrounding skin
- Change the tegaderm prn dislodgement or soiling
- Clean the bladder plate with saline flush if soiled
- The child can be bathed but the bladder plate should not be scrubbed

## Management of the umbilical stump

- Silk suture should be used for the umbilicus
- This is preferred over the umbi clamp to reduce irritation of the bladder mucosa

## Other Considerations

- Provider to provider consultation is available
- If child is stable:
  - No need to transfer until mom can also be discharged
  - Normal diet
  - No antibiotics
  - No activity restrictions

[Return to Prenatal](#)

# Bladder Exstrophy Approval & Citation

Approved by the CSW Bladder Exstrophy team for May 22, 2018 go-live

## CSW Bladder Exstrophy Team:

**Medical Staff Services, Owner**  
**Medical Staff Services, Owner**  
**Ambulatory Urology Nurse**  
**Pharmacy**  
**Orthopedics**  
**Unit Based Educator**  
**Urology**  
**Urology**

Paul Merguerian, MD, MS, FAAP  
Courtney Rowe, MD  
Jean Dearn, RN  
Jia-Jan Seibel, PharmD  
Elizabeth Sheets, ARNP  
Angela Turner, BSN, RN, CPN  
Valerie Nelson, ARNP  
Nicole McMannis, ARNP

## Clinical Effectiveness Team:

**Consultant:**  
**Project Manager:**  
**CE Analyst:**  
**CIS Informatician:**  
**CIS Analyst:**  
**Librarian:**  
**Program Coordinator:**

Jean Popalisky, DNP  
Asa Herrman  
James Johnson  
Carlos Villavicencio, MD, MS/MI  
Julia Hayes, MHIHIM  
Jackie Morton, MLS  
Kristyn Simmons

## Clinical Effectiveness Leadership:

**Medical Director:**  
**Operations Director:**

Darren Migita, MD  
Karen Rancich Demmert, BS, MA

**Retrieval Website:** <http://www.seattlechildrens.org/pdf/bladder-exstrophy-pathway.pdf>

## Please cite as:

Seattle Children's Hospital, Merguerian P, Rowe C, Dearn J, Herrman A, McMannis N, Nelson V, Popalisky J, Sheets E, Turner A, Villavicencio C, 2018 May. Bladder Exstrophy Pathway. Available from: <http://www.seattlechildrens.org/pdf/bladder-exstrophy-pathway.pdf>

# Evidence Ratings

This pathway was developed through local consensus based on published evidence and expert opinion as part of Clinical Standard Work at Seattle Children's. Pathway teams include representatives from Medical, Subspecialty, and/or Surgical Services, Nursing, Pharmacy, Clinical Effectiveness, and other services as appropriate.

When possible, we used the GRADE method of rating evidence quality. Evidence is first assessed as to whether it is from randomized trial or cohort studies. The rating is then adjusted in the following manner (from: Guyatt G et al. J Clin Epidemiol. 2011;4:383-94, Hultcrantz M et al. J Clin Epidemiol. 2017;87:4-13.):

Quality ratings are *downgraded* if studies:

- Have serious limitations
- Have inconsistent results
- If evidence does not directly address clinical questions
- If estimates are imprecise OR
- If it is felt that there is substantial publication bias

Quality ratings are *upgraded* if it is felt that:

- The effect size is large
- If studies are designed in a way that confounding would likely underreport the magnitude of the effect OR
- If a dose-response gradient is evident

## Certainty of Evidence:

★★★★ High: The authors have a lot of confidence that the true effect is similar to the estimated effect

★★★○ Moderate: The authors believe that the true effect is probably close to the estimated effect

★★○○ Low: The true effect might be markedly different from the estimated effect

★○○○ Very low: The true effect is probably markedly different from the estimated effect

Guideline: Recommendation is from a published guideline that used methodology deemed acceptable by the team

Expert Opinion: Based on available evidence that does not meet GRADE criteria (for example, case-control studies).



## Summary of Version Changes

- **Version 1.0 (05/22/2018):** Go live
- **Version 2.0 (2/21/2020):** Clarified Follow-up Coordination section on Post-Operative page

## Medical Disclaimer

Medicine is an ever-changing science. As new research and clinical experience broaden our knowledge, changes in treatment and drug therapy are required.

The authors have checked with sources believed to be reliable in their efforts to provide information that is complete and generally in accord with the standards accepted at the time of publication.

However, in view of the possibility of human error or changes in medical sciences, neither the authors nor Seattle Children's Healthcare System nor any other party who has been involved in the preparation or publication of this work warrants that the information contained herein is in every respect accurate or complete, and they are not responsible for any errors or omissions or for the results obtained from the use of such information.

Readers should confirm the information contained herein with other sources and are encouraged to consult with their health care provider before making any health care decision.

# Bibliography


## Search Methods, *Bladder Exstrophy* Clinical Standard Work

Literature searches were conducted in two phases and executed by a medical librarian, Jackie Morton. The initial search in June 2017, targeted synthesized literature on bladder exstrophy, pelvic fixation methods, the use of anticholinergics, transfusion criteria, as well as the use of antibiotics and pain control and surgical recovery. It was executed in Ovid Medline, Cochrane Database of Systematic Reviews, Embase, National Guideline Clearinghouse and TRIP. The second search, in September 2017, was conducted in Medline and Embase to retrieve primary studies, focusing on pelvic fixation methods, transfusion criteria and the use of anticholinergics or any additional studies on bladder exstrophy. All searches were limited to items published in English, from Jan 2007 to date. The team added 3 citations not retrieved with the search strategy scope and limits. Results were exported to RefWorks for system de-duplication, then to Excel for the screening process.

Jackie Morton, MLS

April 30, 2017

### Identification



Flow diagram adapted from Moher D et al. BMJ 2009;339:bmj.b2535

# Bibliography

Bai Y, Wang X, Li X, et al. Management of catheter-related bladder discomfort in patients who underwent elective surgery. *J Endourol.* 2015;29(6):640-649. Accessed 20150610; 9/20/2017 12:43:24 PM. <https://dx.doi.org/10.1089/end.2014.0670>.

Baradaran N, Cervellione RM, Stec AA, Gearhart JP. Delayed primary repair of bladder exstrophy: Ultimate effect on growth. *J Urol.* 2012;188(6):2336-2341. Accessed 20121112; 9/20/2017 12:43:24 PM. <https://dx.doi.org/10.1016/j.juro.2012.08.037>.

Bryskin RB, Londergan B, Wheatley R, et al. Transversus abdominis plane block versus caudal epidural for lower abdominal surgery in children: A double-blinded randomized controlled trial. *Anesth Analg.* 2015;121(2):471-478. Accessed 9/20/2017 1:03:44 PM. 10.1213/ANE.0000000000000779.

Evaniew N, Khan M, Drew B, Peterson D, Bhandari M, Ghert M. Intrawound vancomycin to prevent infections after spine surgery: A systematic review and meta-analysis. *Eur Spine J.* 2015;24(3):533-542. Accessed 20150319. <https://dx.doi.org/10.1007/s00586-014-3357-0>.

Ferrara F, Dickson AP, Fishwick J, Vashisht R, Khan T, Cervellione RM. Delayed exstrophy repair (DER) does not compromise initial bladder development. *J Pediatr Urol.* 2014;10(3):506-510. Accessed 20140611; 9/20/2017 12:43:24 PM. <https://dx.doi.org/10.1016/j.jpuro.2013.10.026>.

Harper L, Semjen F, Bordes M, et al. Intravesical instillation of ropivacaine reduces bladder spasms following paediatric ureteroneocystostomy. *J Pediatr Urol.* 2007;3(4):301-304. Accessed 9/20/2017 1:03:44 PM. 10.1016/j.jpuro.2006.11.003.

Lako SJ, Steegers MA, van Egmond J, Gardeniers J, Staals LM, van Geffen GJ. Incisional continuous fascia iliaca block provides more effective pain relief and fewer side effects than opioids after pelvic osteotomy in children. *Anesth Analg.* 2009;109(6):1799-1803. Accessed 20091120; 9/20/2017 12:43:24 PM. <https://dx.doi.org/10.1213/ANE.0b013e3181bbc41a>.

# Bibliography

Lo C, Yang SS, Hsieh C, Chang S. Effectiveness of prophylactic antibiotics against post-ureteroscopic lithotripsy infections: Systematic review and meta-analysis. *Surg Infect (Larchmt)*. 2015;16(4):415-420. Accessed 20150725. <https://dx.doi.org/10.1089/sur.2014.013>.

Meara DJ, Livingston NR, Sittitavornwong S, et al. Continuous infusion of bupivacaine for pain control after anterior iliac crest bone grafting for alveolar cleft repair in children. *Cleft Palate Craniofac J*. 2011;48(6):690-694. Accessed 20111102; 9/20/2017 12:43:24 PM. <https://dx.doi.org/10.1597/09-267.1>.

Meyer D, Klarenbeek R, Meyer F. Current concepts in perioperative care for the prevention of deep surgical site infections in elective spinal surgery. *Cen Eur Neurosurg*. 2010;71(3):117-120. Accessed 20100820. <https://dx.doi.org/10.1055/s-0029-1224194>.

New HV, Berryman J, Bolton-Maggs PH, et al. Guidelines on transfusion for fetuses, neonates and older children. *Br J Haematol*. 2016;175(5):784-828. Accessed 11/6/2017 6:21:51 PM. 10.1111/bjh.14233 [doi].

Nguyen TT, Hill S, Austin TM, Whitney GM, Wellons JC, Lam HV. Use of blood-sparing surgical techniques and transfusion algorithms: Association with decreased blood administration in children undergoing primary open craniostylosis repair. *J Neursurg Pediatr*. 2015;16(5):556-563. Accessed 9/20/2017 1:03:44 PM. 10.3171/2015.3.PEDS14663.

Solanki N, Engineer S, Vecham P. Comparison of epidural versus systemic analgesia for major surgeries in neonates and infants. *J Clin Neonatol*. 2017;6(1):23-28. Accessed 9/20/2017 1:03:44 PM. 10.4103/jcn.JCN\_66\_16.

Whyte RK, Jefferies AL, Canadian Paediatric Society, Fetus and Newborn Committee. Red blood cell transfusion in newborn infants. *Paediatr Child Health*. 2014;19(4):213-222. Accessed 11/6/2017 6:21:51 PM.

Wikkelse A, Wetterslev J, Moller AM, Afshari A. Thromboelastography (TEG) or thromboelastometry (ROTEM) to monitor haemostatic treatment versus usual care in adults or children with bleeding. *Cochrane Database of Systematic Reviews*. 2016;(8)-2016 Aug 22. Accessed 20160901; 9/20/2017 12:43:24 PM. <https://dx.doi.org/10.1002/14651858.CD007871.pub3>.

[Return to Home](#)